

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

Pearson Edexcel Level 1/Level 2 GCSE (9–1)

Friday 17 May 2024

Morning (Time: 1 hour 10 minutes)

Paper
reference

1SC0/1CH



Combined Science PAPER 2

Higher Tier

You must have:

Calculator, ruler, Periodic Table (enclosed)

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need*.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must **show all your working out with your answer clearly identified** at the **end of your solution**.

Information

- The total mark for this paper is 60.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question*.
- In questions marked with an **asterisk (*)**, marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P74438A

©2024 Pearson Education Ltd.
F:1/1/1/1/1/



Pearson

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box . If you change your mind about an answer, put a line through the box and then mark your new answer with a cross .

1 Barium hydroxide reacts with dilute hydrochloric acid to form barium chloride solution and water.

(a) (i) Complete the balanced equation for the reaction by adding a **number** in front of HCl(aq). (1)



(ii) State what you would **see** during the reaction. (1)

.....
.....
.....
.....
.....

(b) A student investigated how the pH of the mixture changed as barium hydroxide was added to dilute hydrochloric acid.

The student used this method.

step 1 measure out 50 cm³ of dilute hydrochloric acid into a beaker using a measuring cylinder

step 2 use a glass rod to place a drop of the acid onto a piece of universal indicator paper and record the pH

step 3 add one spatula measure of barium hydroxide to the acid in the beaker and stir

step 4 use the glass rod to place a drop of the mixture onto a new piece of universal indicator paper and record the pH again

step 5 repeat steps 3 and 4 until there is no further change in the pH.



(i) Name a piece of equipment that could be used to measure the pH of a substance more accurately than universal indicator paper.

(1)

(ii) Explain why, in step 3, the mixture was stirred after adding the barium hydroxide.

(2)

(iii) Figure 1 shows the student's results.


number of spatula measures of barium hydroxide	pH of mixture
0	1
1	1
2	1
3	1
4	3
5	8
6	12
7	13
8	13

Figure 1



Plot a graph of the pH of the mixture against the number of spatula measures of barium hydroxide.

(3)



(iv) Use the graph to find the pH of the mixture when 4.5 spatula measures of barium hydroxide are added.

(1)

pH of the mixture =

(Total for Question 1 = 9 marks)



P 7 4 4 3 8 A 0 5 2 0

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



2 Sodium carbonate has the formula Na_2CO_3 .

(a) Sodium carbonate contains Na^+ ions and CO_3^{2-} ions.

(i) The atomic number of sodium is 11.

What is the electronic configuration of the Na^+ ion?

(1)

- A** 1
- B** 2.8
- C** 2.8.1
- D** 2.8.2

(ii) Explain why solid sodium carbonate **cannot** conduct electricity but a solution of sodium carbonate **can** conduct electricity.

(3)

(b) Calculate the percentage by mass of sodium in sodium carbonate, Na_2CO_3 .

$$\text{percentage by mass of element} = \frac{\text{total relative atomic mass of element}}{\text{relative formula mass of compound}} \times 100$$

(relative atomic masses: C = 12, O = 16, Na = 23)

(3)

percentage by mass of sodium =

(Total for Question 2 = 7 marks)



3 This question is about the extraction of metals.

(a) Give **two** advantages of obtaining metals by recycling rather than by extracting them from their metal ores.


(2)

1

2

(b) (i) Small amounts of some metals are extracted using displacement reactions.

In one process, zinc dust is used to precipitate cadmium metal from a solution containing cadmium ions.



Explain why this displacement reaction can be described as a **redox reaction**.

(3)



(ii) The formula of the cadmium ion is Cd^{2+} .
The formula of the phosphate ion is PO_4^{3-} .

Which is the formula of cadmium phosphate?

(1)

- A $\text{Cd}_2(\text{PO}_4)_3$
- B $\text{Cd}_3\text{PO}_{12}$
- C $\text{Cd}_3(\text{PO}_4)_2$
- D $\text{Cd}_3\text{P}_2\text{O}_8$

(c) One of the alternative biological methods of extracting metals from very low-grade ores is bioleaching using bacteria.

Give one **disadvantage** of this method of extracting metals from low-grade ores.

(1)

(d) Lead is low in the reactivity series.

Describe how to obtain a sample of lead from some lead oxide in the laboratory.

(2)

(Total for Question 3 = 9 marks)



P 7 4 4 3 8 A 0 9 2 0

4 (a) Water, acidified with dilute sulfuric acid, was electrolysed for 10 minutes using inert electrodes.

Figure 2 shows the apparatus used.




Figure 2

(i) In this electrolysis, the acidified water is an electrolyte.





Explain why acidified water is an electrolyte.

(2)

(ii) Hydrogen collects at the negative electrode and oxygen collects at the positive electrode.

Which of these shows the results after 10 minutes of this electrolysis?

(1)



(iii) Complete and balance the half equation for the formation of oxygen at the positive electrode in this electrolysis.

(2)



(b) Copper sulfate solution was electrolysed for 10 minutes using copper electrodes.

Figure 3 shows the mass of the cathode and the appearance of the copper sulfate solution before electrolysis and after electrolysis.

	mass of cathode in g	appearance of copper sulfate solution
before electrolysis	5.32	pale blue solution
after electrolysis	5.87	pale blue solution

Figure 3

(i) Describe what should be done to the copper cathode, after it has been removed from the copper sulfate solution, before its final mass is determined.

(2)

(ii) Explain, in terms of ions, the change in mass of the cathode shown in Figure 3.

(2)



(iii) Explain why the appearance of the copper sulfate solution did not change during the electrolysis.

(2)

(Total for Question 4 = 11 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



P 7 4 4 3 8 A 0 1 3 2 0

5 This question is about the properties of different substances.

(a) Silicon tetrachloride is a simple molecular covalent compound.

(i) A molecule of silicon tetrachloride is composed of a silicon atom and four chlorine atoms.

- a silicon atom has 4 outer electrons
- a chlorine atom has 7 outer electrons

Draw a dot and cross diagram of a molecule of silicon tetrachloride, SiCl_4 .

Show outer electrons only.

(2)

(ii) Explain why simple molecular covalent compounds such as silicon tetrachloride have low melting and boiling points.

(2)



(b) Part of the structure of rubidium bromide is shown in Figure 4.




Figure 4

Which row shows the most likely melting point and boiling point of rubidium bromide?

(1)

	melting point in °C	boiling point in °C
<input type="checkbox"/> A	6.93	134.0
<input type="checkbox"/> B	69.3	134.0
<input type="checkbox"/> C	69.3	1340
<input type="checkbox"/> D	693	1340



***c) Diamond and graphite are two forms of carbon.**

Figure 5 shows how the carbon atoms are arranged in a part of the structure of each of these forms of carbon.




Figure 5

- diamond is one of the hardest known substances on Earth and is used in cutting tools.
- graphite is soft and flaky.
- diamond is a poor electrical conductor, but graphite is a good electrical conductor.

Explain, in terms of structure and bonding, these properties of diamond and graphite.

(6)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 5 = 11 marks)



P 7 4 4 3 8 A 0 1 7 2 0

6 (a) An investigation was carried out on the reactivity of four metals, **D**, **E**, **F** and **G**.

Equal sized pieces of these metals were placed in excess dilute hydrochloric acid and left for three minutes.

Figure 6 shows the observations of the reactions for metals **D**, **E** and **F**.

metal	observations with dilute hydrochloric acid
D	Bubbles formed quickly. After three minutes all the metal had reacted.
E	Bubbles formed very quickly. No metal remaining after three minutes.
F	A few bubbles were seen to form. The metal looked unchanged after three minutes.
G	

Figure 6

The order of reactivity for these metals is shown in Figure 7.

E	D	G	F
most reactive		least reactive	

Figure 7

(i) Use the information in Figure 6 and Figure 7 to suggest the observations that would be made for metal **G**.

(2)

.....

.....

.....



(ii) The dilute hydrochloric acid used in this reaction is a strong acid.

Explain the meaning of the terms **dilute** and **strong acid**.

(4)

dilute

.....
.....
.....

strong acid

.....
.....
.....

(b) The formula of lead ethanoate is $\text{Pb}(\text{CH}_3\text{COO})_2$.

Calculate the number of **atoms** that combine together to form 16.25 g of lead ethanoate.

(relative atomic masses: H = 1.00, C = 12.0, O = 16.0, Pb = 207)

Avogadro number = 6.02×10^{23})

(4)

number of atoms =



P 7 4 4 3 8 A 0 1 9 2 0

(c) Iron is more reactive than copper.

Iron will displace copper from copper nitrate solution.

Two possible balanced equations for the reaction are



It was found that 2.24 g of iron reacted with excess copper nitrate solution to form 3.81 g of copper.

Carry out a calculation, using the information above, to show which equation represents the reaction taking place.

(relative atomic masses: Fe = 56.0, Cu = 63.5)

(3)

(Total for Question 6 = 13 marks)

TOTAL FOR PAPER = 60 MARKS



Pearson Edexcel Level 1/Level 2 GCSE (9–1)

Friday 17 May 2024

**Paper
reference**

1SC0/1CH



Combined Science

PAPER 2

Higher Tier

Periodic Table Insert

Do not return this Insert with the question paper

P74438A

©2024 Pearson Education Ltd.
F:1/1/1/1/1/



Turn over ►



Pearson

The periodic table of the elements

1	2											
7 Li lithium 3	9 Be beryllium 4											
23 Na sodium 11	24 Mg magnesium 12											
39 K potassium 19	40 Ca calcium 20	45 Sc scandium 21	48 Ti titanium 22	51 V vanadium 23	52 Cr chromium 24	55 Mn manganese 25	56 Fe iron 26	59 Co cobalt 27	59 Ni nickel 28	63.5 Cu copper 29	65 Zn zinc 30	
85 Rb rubidium 37	88 Sr strontium 38	89 Y yttrium 39	91 Nb niobium 41	93 Zr zirconium 40	96 Mo molybdenum 42	[98] Tc technetium 43	101 Ru ruthenium 44	103 Rh rhodium 45	106 Pd palladium 46	108 Ag silver 47	112 Cd cadmium 48	115 In indium 49
133 Cs caesium 55	137 Ba barium 56	139 La[*] lanthanum 57	178 Hf hafnium 72	181 Ta tantalum 73	184 W tungsten 74	186 Re rhenium 75	190 Os osmium 76	192 Ir iridium 77	195 Pt platinum 78	197 Au gold 79	201 Hg mercury 80	204 Tl thallium 81
Key <div style="border: 1px solid black; padding: 5px; display: inline-block;"> relative atomic mass atomic symbol <small>name</small> atomic (proton) number </div>												
11 B boron 5	12 C carbon 6	14 N nitrogen 7	16 O oxygen 8	19 F fluorine 9	20 Ne neon 10	27 Al aluminium 13	28 Si silicon 14	31 P phosphorus 15	32 S sulfur 16	35.5 Cl chlorine 17	40 Ar argon 18	
1 H hydrogen 1	3 He helium 2	4 He helium 2	5	6	7	0						

* The elements with atomic numbers from 58 to 71 are omitted from this part of the periodic table.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.

